

# Problem Set 3: Polynomial Approximation

July 26, 2023

**Problem 1** (Polynomial approximation of monomials). First, compute the Chebyshev coefficients of the monomial  $m^{(n)}(x) = x^n$ . (Doing this via  $T_k(\frac{1}{2}(z + z^{-1})) = \frac{1}{2}(z^k + z^{-k})$  formulation may be easiest.) How small can  $k$  be such that the Chebyshev truncation  $m_k^{(n)}$  a good approximation of  $m^{(n)}$ :

$$\|m^{(n)} - m_k^{(n)}\|_{[-1,1]} \leq \varepsilon?$$

**Problem 2** (Chebyshev interpolation [Tre19]). The *Chebyshev interpolant* of a function  $f$ , denoted  $p_d$ , is the unique degree- $d$  polynomial such that  $p_d(x_j) = f(x_j)$  for all  $x_j = \cos(j\pi/d)$ ,  $j = 0, 1, \dots, d$ . Prove that<sup>1</sup>

$$\|f(x) - p_d(x)\|_{[-1,1]} \leq 2 \sum_{\ell \geq d} |a_\ell|.$$

Hint: when is  $T_k(x_j) = T_\ell(x_j)$  for all points  $\{x_j\}$ ?

**Problem 3** (Jackson theorems, [Tre19]). Let  $f : [-1, 1] \rightarrow \mathbb{R}$  be absolutely continuous and suppose  $f$  is of bounded variation, meaning that  $\int_{-1}^1 |f'(x)| dx \leq V$ . Then show that the Chebyshev coefficients of  $f$  satisfy

$$|a_k| \leq \frac{2V}{\pi k}.$$

**Problem 4** (Optimal polynomial approximations; upper and lower bounds). Consider a function  $f : [-1, 1] \rightarrow \mathbb{R}$  with a Chebyshev expansion  $f(x) = \sum_{k \geq 0} a_k T_k(x)$ . Prove that

$$\left(\frac{1}{2} \sum_{k=n+1}^{\infty} a_k^2\right)^{\frac{1}{2}} \leq \min_{\substack{p \in \mathbb{R}[x] \\ \deg p = n}} \|f(x) - p(x)\|_{[-1,1]} \leq \sum_{k=n+1}^{\infty} |a_k|$$

For what kind of Chebyshev coefficient decay is this characterization tight up to constants?

## References

- [Tre19] Lloyd N. Trefethen. *Approximation theory and approximation practice, extended edition*. Extended edition [of 3012510]. Philadelphia, PA: Society for Industrial and Applied Mathematics, 2019, pp. xi+363. ISBN: 978-1-611975-93-2. DOI: [10.1137/1.9781611975949](https://doi.org/10.1137/1.9781611975949) (page 1).

---

<sup>1</sup>Recall that our approximation results used that  $\|f(x) - f_d(x)\|_{[-1,1]} \leq \sum_{\ell \geq d} |a_\ell|$ . So, Chebyshev interpolants  $p_d$  give the same results as Chebyshev truncations  $f_d$ , up to a constant factor. Interpolants have the advantage of being computable in  $d + 1$  function evaluations.